



## **Antioxidant Activity of Fish Protein Hydrolysates in in vitro Assays and in Oil-in-Water Emulsions.**

**Farvin, Sabeena; Andersen, Lisa Lystbæk; Jacobsen, Charlotte; Nielsen, Henrik Hauch; Jessen, Flemming**

*Publication date:*  
2011

[Link back to DTU Orbit](#)

*Citation (APA):*  
Farvin, S., Andersen, L. L., Jacobsen, C., Nielsen, H. H., & Jessen, F. (2011). *Antioxidant Activity of Fish Protein Hydrolysates in in vitro Assays and in Oil-in-Water Emulsions..* Abstract from 102nd AOCS Annual Meeting & Expo, Cincinnati, Ohio, United States.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Farvin KHS., Andersen LL., Jacobsen C., Nielsen HH., Jessen F.**

**Antioxidant activity of fish protein hydrolysates in *in vitro* assays and in oil-in-water emulsions.**

The aim of this study was to screen different protein hydrolysates with respect to their antioxidative properties in order to select the most promising extracts for further evaluation in oil-in-water emulsions. Three fractions of protein hydrolysates (Crude, >5kDa and <5kDa) from cod and plaice were screened for antioxidant activity by using four *in vitro* assays viz, 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity, reducing power, ferrous iron-chelating and inhibition of lipid oxidation in liposome model system. Cod protein hydrolysates exhibited significantly higher antioxidative effect in various antioxidant assays and were selected for further study. The lower fraction <5kDa was further fractionated by using a 3kDa cut-off membrane to get two fractions 3-5kDa and <3kDa. Thus, four fractions of cod protein hydrolysates (Crude, >5kDa, 3-5kDa and <3kDa) were further screened by using the above mentioned antioxidant assays. Interestingly, it was found that the fraction <3kDa showed very good reducing power, Fe<sup>2+</sup> chelating activity and DPPH radical scavenging activity. Moreover, the 3-5kDa fraction was very effective in retarding lipid oxidation in liposome model system. These two fractions were therefore selected for further evaluation in 5% fish oil in water emulsions.